

Summer 7-2018

Using Synthetic Lethality to Sensitize Pediatric Brain Cancer Cells to the DNA Damaging Effects of Radiation Therapy

Megan Ruckman

megan.ruckman@huskers.unl.edu

Rose Nelson

rkan41898@gmail.com

Forrest Kievit

fkievit2@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/ucareresearch>



Part of the [Other Biomedical Engineering and Bioengineering Commons](#)

Ruckman, Megan; Nelson, Rose; and Kievit, Forrest, "Using Synthetic Lethality to Sensitize Pediatric Brain Cancer Cells to the DNA Damaging Effects of Radiation Therapy" (2018). *UCARE Research Products*. 158.

<https://digitalcommons.unl.edu/ucareresearch/158>

This Poster is brought to you for free and open access by the UCARE: Undergraduate Creative Activities & Research Experiences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in UCARE Research Products by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Using Synthetic Lethality to Sensitize Pediatric Brain Cancer Cells to the DNA Damaging Effects of Radiation Therapy

Rose Nelson, Megan Ruckman, & Dr. Forrest Kievit
Department of Biological Systems Engineering
University of Nebraska-Lincoln

Nebraska
Engineering




INTRODUCTION

Brain tumors are the leading cause of cancer related deaths in children. As a treatment, radiotherapy is very effective, but it also comes with many severe side effects, such as radiation-induced adverse developmental and psychosocial effects.² Cancer, because it is a mutagenic disease, is by definition a disease of DNA repair pathways. Here we utilize drugs to inhibit various DNA repair pathways with the purpose of sensitizing cancer cells to radiotherapy so that lower doses may be used. Hopefully this would improve the lifelong outcomes of patients that would otherwise not survive or would be left with lifelong neurocognitive problems.

METHODS

AlamarBlue was used to determine the number of cells in each well by quantifying metabolic activity of cultured cells. The metabolically active cells reduce the blue AlamarBlue substrate into a pink product (Fig. 1a). A measure of cell kill or cell viability is determined by comparing the conversion of AlamarBlue to a standardized curve (Fig. 1b). Hydrogen peroxide was used to mimic the DNA damaging effects of radiotherapy by causing single and double stranded breaks (Fig. 3a). Various drugs were used to inhibit the repair pathways of the cells (Fig. 3b-f). Four different cell lines underwent these same experimental conditions with each inhibitor drug (Fig. 1c).



RESULTS




Figure 3. a) Cell viability of the different cell lines after being treated with the DNA-repair inhibiting drugs at different concentrations of hydrogen peroxide. AlamarBlue was used to measure the viability of the cells each day for three days after exposure to hydrogen peroxide, b) LD₅₀ of hydrogen peroxide of control and when treated with each of the four drugs at each day, c) microscope image of U118 cell line, d) microscope image of C6 cell line, e) microscope image of RES 196 cell line, f) microscope image of H59EP cell line.

THE INHIBITOR DRUGS





Figure 2. a) Hydrogen peroxide treatment mimics the DNA damage caused by ionizing radiation by generating single-strand and double-strand breaks. b) E3330 targets APE1/Ref-1 and is a redox inhibitor of the protein by destabilizing the protein's structure. APE1/Ref-1 is a protein that controls both DNA repair and redox reactions that allow cancer cells to thrive.³ c) KU-55933 inhibits the ataxia-telangiectasia mutated (ATM), which is the activated by DNA double stranded breaks and initiates signaling cascades that start DNA damage response.⁴ d) AZD2461 inhibits the healing of single-stranded breaks by binding to the nuclear enzyme poly(ADP-ribose) polymerase (PARP) a protein that plays a significant role in DNA repair,⁵ e) NU7026 is an inhibitor of DNA-dependent protein kinase (DNA-PK), a complex that acts as a scaffold for DNA repair, and inhibits repair of double stranded breaks in cells that have the DNA-PK. NU7026 has been shown to impair cell survival.⁶ f) molecular structure of the drugs used in this experiment.

CYTATION

The Cytation imager will be used in the Fall for experiments. The cells get stained with Hoechst stain, which stains the nuclei of each cell. The Cytation imager counts the actual number of cells by counting the stained nuclei instead of measuring metabolic activity, like AlamarBlue (Fig. 4a). This experiment shows that there is a linear correlation between number of cells in the well and cell counts provided by the Cytation (Fig. 4b).



DISCUSSION

The cancer cells saw a high toxicity with the concentrations of AZD2461 and E3330 which may have lead to some of the inconclusive data that was recovered (Fig. 2a). Lower concentrations of these two drugs will be used in future experiments, this should increase the accuracy of the LD₅₀ of hydrogen peroxide. We also saw an increase in the LD₅₀ of many of the cell lines when treated with NU7026 and KU55933, which may indicate that knocking out the one repair pathway actually allowed the cells to thrive rather than die (Fig. 2b). The statistical significance of this data must still be determined and more experiments must be done to provide more conclusive information. In conclusion, there may be a correlation between certain drug treatments and increased sensitivity to DNA damage similar to that caused by radiotherapy. However, more research must be done to draw more reliable conclusions.

ACKNOWLEDGEMENTS

Special Thanks to the UCARE program, as well as the Buffett Cancer Center and Pediatric Cancer Research Group at UNMC for a seed grant that helped fund this work.

REFERENCES

1. Khalil, HS et al. "Pharmacological inhibition of ATM by KU55933 stimulates ATM transcription." *Experimental Biology and Medicine*. 237 (2012) 622-634.
2. Kievit, Forrest M. et al. "Nanoparticle mediated silencing of DNA repair sensitizes pediatric brain tumor cells to γ -irradiation." *Molecular Oncology* 9 (2015) 1071-1080.
3. Laev SS. et al. "Inhibitors of nuclease and redox activity of apurinic/aprimidinic endonuclease 1/redox effector (APE1/Ref-1)." *Bioorganic & Medicinal Chemistry*. 25.9 (2017) 2531-2544.
4. Martin, Purschke, et al. "Phototoxicity of Hoechst 33342 in Time-Lapse Fluorescence Microscopy." *Photochemical & Photobiological Sciences*, vol. 9, no. 12, Dec. 2010, pp. 1634-1639.
5. Niazi, MT. et al. "Effects of DNA-dependent protein kinase inhibition by NU7026 on DNA repair and cell survival in irradiated gastric cancer cell line N87." *Current Oncology*. 21.2 (2014) 91-96.
6. O'Connor, L.O. et al. "The PARP inhibitor AZD2461 provides insights into the role of PARP3 inhibition for both synthetic lethality and tolerability with chemotherapy in preclinical models." *Cancer Res*. 76 (2016) 6084-6094.